

THE 2019 POSITIVELY AWARE HIV DRUG CHART



Each drug is recommended for use in combination with other antiretrovirals (except any single-tablet regimen, including Atripla, Biktarvy, Complera, Delstrigo, Genvoya, Juluca, Odefsey, Stribild, Symfi, Symfi Lo, Symtuza, and Triumeq). For a complete description of DHHS drug recommendations, go to aidsinfo.nih.gov/guidelines. Consult with a physician about which drug therapy and dose is appropriate for you; see drug page or package insert for complete information. Drug chart information is current as of February 1, 2019. Look up your drug by adding its name after you type positivelyaware.com/ into your browser (for example, positivelyaware.com/triumeq).

These medications contain a complete regimen all in one:

SINGLE TABLET REGIMENS (Multiple drug classes; see pregnancy information for integrase inhibitors below.)

Atripla ✓ RECOMMENDED INITIAL REGIMEN IN CERTAIN CLINICAL SITUATIONS

efavirenz/emtricitabine/tenofovir DF (EFV/FTC/TDF)
One tablet (600 mg efavirenz/200 mg emtricitabine/300 mg tenofovir DF), once daily. Take on an empty stomach, preferably at bedtime.

Complera ✓ RECOMMENDED INITIAL REGIMEN IN CERTAIN CLINICAL SITUATIONS

rilpivirine/emtricitabine/tenofovir DF (RPV/FTC/TDF)
One tablet (25 mg rilpivirine/200 mg emtricitabine/300 mg tenofovir DF), once daily. Take with a meal.

Genvoya ✓ RECOMMENDED INITIAL REGIMEN IN CERTAIN CLINICAL SITUATIONS

elvitegravir/cobicistat/emtricitabine/tenofovir alafenamide (EVG/COBI/FTC/TAF)
One tablet (150 mg elvitegravir/150 mg cobicistat/200 mg emtricitabine/10 mg tenofovir alafenamide), once daily. Take with food.

Odefsey ✓ RECOMMENDED INITIAL REGIMEN IN CERTAIN CLINICAL SITUATIONS

rilpivirine/emtricitabine/tenofovir alafenamide (RPV/FTC/TAF)
One tablet (25 mg rilpivirine/200 mg emtricitabine/25 mg tenofovir alafenamide), once daily. Take with a meal.

Symfi ✓ RECOMMENDED INITIAL REGIMEN IN CERTAIN CLINICAL SITUATIONS

efavirenz/lamivudine/tenofovir DF (EFV/3TC/TDF)
One tablet (600 mg efavirenz/300 mg lamivudine/300 mg tenofovir DF), once daily. Should be taken on an empty stomach, preferably at bedtime.

Symtuza ✓ RECOMMENDED INITIAL REGIMEN IN CERTAIN CLINICAL SITUATIONS

darunavir/cobicistat/emtricitabine/tenofovir alafenamide (DRV/COBI/FTC/TAF)
One tablet (800 mg darunavir/150 mg cobicistat/200 mg emtricitabine/10 mg tenofovir alafenamide), once daily for patients without darunavir- or tenofovir-related drug resistance. Take with food.

dolutegravir/lamivudine ✓ INITIAL REGIMEN TO CONSIDER WHEN ABC, TAF, AND TDF CANNOT BE USED OR ARE NOT OPTIMAL

(DTG/3TC)
NOT YET APPROVED AT PRESS TIME.
One tablet, once daily. Tablet contains 50 mg of the INSTI dolutegravir plus 300 mg of the NRTI lamivudine. Lamivudine dosing needs to be adjusted for adults and children who have decreased kidney function (creatinine clearance less than 50 mL/min). Take with or without food.

PHOTO UNAVAILABLE

Biktarvy ★ RECOMMENDED INITIAL REGIMEN FOR MOST PEOPLE

bictegravir/emtricitabine/tenofovir alafenamide (BIC/FTC/TAF)
One tablet (50 mg bictegravir/200 mg emtricitabine/25 mg tenofovir alafenamide), once daily. For adults who are treatment-naïve, or on treatment and virologically suppressed for at least 3 months, with no history of treatment failure or resistance mutations associated with components of Biktarvy. Take with or without food.

Delstrigo ✓ RECOMMENDED INITIAL REGIMEN IN CERTAIN CLINICAL SITUATIONS

doravirine/lamivudine/tenofovir DF (DOR/3TC/TDF)
One tablet (100 mg doravirine/300 mg lamivudine/300 mg tenofovir DF), once daily. Take with or without food.

Juluca ✓ RECOMMENDED AS CONTINUATION THERAPY FOR PEOPLE WITH UNDETECTABLE HIV VIRAL LOAD FOR AT LEAST 6 MONTHS

dolutegravir/rilpivirine (DTG/RPV)
One tablet (50 mg dolutegravir/25 mg rilpivirine), once daily. For adults who are virologically suppressed on a current ART regimen for at least 6 months and who have no history of treatment failure or resistance mutations associated with rilpivirine or dolutegravir. Take with a meal.

Stribild ✓ RECOMMENDED INITIAL REGIMEN IN CERTAIN CLINICAL SITUATIONS

elvitegravir/cobicistat/emtricitabine/tenofovir DF (EVG/COBI/FTC/TDF)
One tablet (150 mg elvitegravir/150 mg cobicistat/200 mg emtricitabine/300 mg tenofovir DF), once daily. Take with food.

Symfi Lo ✓ RECOMMENDED INITIAL REGIMEN IN CERTAIN CLINICAL SITUATIONS

efavirenz/lamivudine/tenofovir DF (EFV/3TC/TDF)
One tablet (400 mg efavirenz/300 mg lamivudine/300 mg tenofovir DF), once daily. Should be taken on an empty stomach, preferably at bedtime.

Triumeq ★ RECOMMENDED INITIAL REGIMEN FOR MOST PEOPLE IF HLA-B*5701 NEGATIVE

dolutegravir/abacavir/lamivudine (DTG/ABC/3TC)
One tablet (50 mg dolutegravir/600 mg abacavir/300 mg lamivudine), once daily. One additional 50 mg tablet of Tricyclic 12 hours apart in people with viral resistance to INSTIs or when taken with certain other medications. Take with or without food.

LONG ACTING REGIMEN

cabotegravir LA/rilpivirine LA ● DHHS RECOMMENDATION NOT YET ESTABLISHED

(CAB LA/RPV LA)
NOT YET APPROVED AT PRESS TIME.
Long-acting injection of 400 mg cabotegravir plus 600 mg rilpivirine every 4 or 8 weeks in studies. Dose in studies consisted of two 2 mL injections. Rilpivirine must be taken with food. Induction phase with oral medication used in research and will be used when approved. Cabotegravir lead-in dose tablet may not otherwise be available on the market.

PHOTO UNAVAILABLE

These medications must be taken in combination with other HIV meds:

INTEGRASE INHIBITORS (Pregnancy testing should be performed in those of childbearing potential prior to initiation.)

Isentress HD ★ RECOMMENDED AS COMPONENT OF INITIAL REGIMEN FOR MOST PEOPLE

and Isentress raltegravir (RAL)
Isentress HD: Two 600 mg film-coated tablets, once daily.
Isentress: One 400 mg film-coated tablet, twice daily. Take with or without food.

Tivicay ★ RECOMMENDED AS COMPONENT OF INITIAL REGIMEN FOR MOST PEOPLE

dolutegravir (DTG)
One 50 mg tablet, once daily for people on HIV therapy for the first time or treatment-experienced without previous INSTI resistance. Twice-daily dosing for people who have viral resistance to INSTIs or when taken with certain other medications. Take with or without food.

PROTEASE INHIBITORS (Boosted and unboosted)

Evotaz ✓ RECOMMENDED AS COMPONENT OF INITIAL REGIMEN IN CERTAIN CLINICAL SITUATIONS

atazanavir/cobicistat (ATV/COBI)
One tablet (300 mg atazanavir/150 mg cobicistat), once daily. Take with food.

Prezcobix ✓ RECOMMENDED AS COMPONENT OF INITIAL REGIMEN IN CERTAIN CLINICAL SITUATIONS

darunavir/cobicistat (DRV/COBI)
One tablet (800 mg darunavir/150 mg cobicistat), once daily in patients with no darunavir drug resistance. Take with food.

Prezista ✓ RECOMMENDED AS COMPONENT OF INITIAL REGIMEN IN CERTAIN CLINICAL SITUATIONS

darunavir (DRV)
One 800 mg tablet with 100 mg Norvir or 150 mg Tybost, once daily for first-time therapy and treatment-experienced adults without Prezista-related resistance; or one 600 mg tablet plus 100 mg Norvir, twice daily for treatment-experienced people with Prezista-related resistance. Prezista must be taken with Norvir or Tybost. Take with food.

Reyataz ✓ RECOMMENDED AS COMPONENT OF INITIAL REGIMEN IN CERTAIN CLINICAL SITUATIONS

atazanavir sulfate (ATV)
One 300 mg capsule plus 100 mg Norvir or 150 mg Tybost, once daily; or two 200 mg capsules (without boosting), once daily for treatment-naïve adults (see the POSITIVELY AWARE HIV Drug Guide for details). Take with food. **GENERIC IS AVAILABLE.**

PK ENHANCERS (Boosters)

Tybost ✓ USED ONLY AS A BOOSTER FOR OTHER DRUGS; RECOMMENDED AS COMPONENT OF INITIAL REGIMEN IN CERTAIN CLINICAL SITUATIONS

cobicistat (COBI)
Used only for boosting—not an antiretroviral. 150 mg once daily taken at the same time with either Prezista 800 mg or Reyataz 300 mg. Take with food.

Norvir ✓ USED ONLY AS A BOOSTER FOR OTHER DRUGS; RECOMMENDED AS COMPONENT OF INITIAL REGIMEN IN CERTAIN CLINICAL SITUATIONS

ritonavir (RTV)
Used only for boosting, also an antiretroviral. 100–200 mg, dosed once or twice daily with another PI. Must be taken with food. **GENERIC IS AVAILABLE.**

NUCLEOSIDE REVERSE TRANSCRIPTASE INHIBITORS ("Nukes") ★ All "nukes" are recommended.

Cimduo RECOMMENDED FOR INITIAL ART FOR MOST PERSONS WHEN COMBINED WITH TIVICAY OR ISENTRESS

lamivudine/tenofovir DF (3TC/TDF)
One tablet (300 mg lamivudine/300 mg tenofovir DF), once daily, taken in combination with other ARVs. Take with or without food.

Descovy RECOMMENDED AS A COMPONENT OF INITIAL REGIMEN FOR MOST PEOPLE

emtricitabine/tenofovir alafenamide (FTC/TAF)
One tablet (200 mg emtricitabine/25 mg tenofovir alafenamide), once daily. Take with or without food.

Emtriva RECOMMENDED AS A COMPONENT OF INITIAL REGIMEN FOR MOST PEOPLE

emtricitabine (FTC)
One 200 mg capsule, once daily. Take with or without food.

Epivir RECOMMENDED AS COMPONENT OF INITIAL REGIMEN FOR MOST PEOPLE WHEN USED IN COMBINATION WITH DOLUTEGRAVIR AND ABACAVIR

lamivudine (3TC)
One 300 mg tablet, once daily; or one 150 mg tablet, twice daily. Take with or without food. **GENERIC IS AVAILABLE.**

Epzicom RECOMMENDED AS COMPONENT OF INITIAL REGIMEN FOR MOST PEOPLE IF HLA-B*5701-NEGATIVE WHEN USED IN COMBINATION WITH DOLUTEGRAVIR (AS TRIUMEQ)

abacavir/lamivudine (ABC/3TC)
One tablet (600 mg abacavir/300 mg lamivudine), once daily. Take with or without food. **GENERIC IS AVAILABLE.**

Truvada RECOMMENDED AS A COMPONENT OF INITIAL REGIMEN FOR MOST PEOPLE

emtricitabine/tenofovir DF (FTC/TDF)
One tablet (200 mg emtricitabine/300 mg tenofovir DF), once daily. Take with or without food. **APPROVED AS GENERIC, WHICH IS NOT YET COMMERCIALY AVAILABLE.**

Viread RECOMMENDED AS A COMPONENT OF INITIAL REGIMEN FOR MOST PEOPLE

tenofovir disoproxil fumarate (TDF)
One 300 mg tablet, once daily. Take with or without food. **GENERIC IS AVAILABLE.**

Ziagen RECOMMENDED AS COMPONENT OF INITIAL REGIMEN FOR MOST PEOPLE IF HLA-B*5701-NEGATIVE WHEN USED IN COMBINATION WITH DOLUTEGRAVIR AND LAMIVUDINE (AS TRIUMEQ)

abacavir sulfate (ABC)
Two 300 mg tablets, once daily; or one 300 mg tablet, twice daily. Take with or without food. **GENERIC IS AVAILABLE.**

NON NUCLEOSIDE REVERSE TRANSCRIPTASE INHIBITORS (Non nukes")

Edurant ✓ RECOMMENDED AS COMPONENT OF INITIAL REGIMEN IN CERTAIN CLINICAL SITUATIONS IN COMBINATION WITH DESCOVY OR TRUVADA (AS ODEFSEY OR COMPLERA)

rilpivirine (RPV)
One 25 mg tablet, once daily. Take with a meal.

Intelence ✓ FOR TREATMENT-EXPERIENCED PATIENTS WITH VIRAL STRAINS RESISTANT TO AN NRTI AND OTHER ANTIRETROVIRAL DRUGS ONLY

etravirine (ETR)
One 200 mg tablet, twice daily. Take with food.

Pifeltro ✓ RECOMMENDED AS COMPONENT OF INITIAL REGIMEN IN CERTAIN CLINICAL SITUATIONS IN COMBINATION WITH DESCOVY OR TRUVADA, OR WITH CIMDUO (AS DELSTRIGO)

doravirine (DOR)
One 100 mg tablet, once daily. Take with or without food.

Sustiva ✓ RECOMMENDED AS COMPONENT OF INITIAL REGIMEN IN CERTAIN CLINICAL SITUATIONS (AS A PART OF ATRIPLA, SYMFI, AND SYMFI LO, OR WITH DESCOVY OR TRUVADA)

efavirenz (EFV)
One 600 mg tablet, once daily. Take on an empty stomach, preferably at bedtime. **GENERIC IS AVAILABLE.**

ENTRY/ATTACHMENT INHIBITORS

Selzentry maraviroc (MVC)

▼ NOT RECOMMENDED AS COMPONENT OF AN INITIAL REGIMEN
ENTRY INHIBITOR: CCR5 ANTAGONIST
150, 300, or 600 mg (available in 150 and 300 mg tablets), twice daily, depends on other medications used (see the POSITIVELY AWARE HIV Drug Guide for details). Take with or without food.

Trogarzo ibalizumab-uiyk (IBA)

▼ FOR PEOPLE LACKING SUFFICIENT TREATMENT OPTIONS
CD4 POST-ATTACHMENT INHIBITOR
Administered once every two weeks via intravenous infusion. Treatment begins with an IV loading (starting) dose of 2,000 mg, followed by an 800 mg IV infusion maintenance dose given every two weeks thereafter.

fostemsavir (FTR)

● DHHS RECOMMENDATION NOT YET ESTABLISHED
ATTACHMENT INHIBITOR
NOT YET APPROVED AT PRESS TIME.
In clinical trials, the investigational dose taken forward for further study was 600 mg, sometimes once daily and sometimes twice daily. Doses were taken after eating. For highly-treatment experienced individuals with history of antiretroviral resistance.

LEGACY DRUGS The following drugs are no longer or rarely prescribed:

 **Aptivus**
tipranavir (TPV)

 **Combivir**
lamivudine/zidovudine (3TC/AZT)

 **Crixivan**
indinavir (IDV)

 **Fuzeon**
enfuvirtide (T-20, or ENF)

 **Invirase**
saquinavir (SQV)

 **Kaletra**
lopinavir/ritonavir (LPV/r)

 **Lexiva**
fosamivir calcium (FPV)

 **Rescriptor**
delavirdine (DLV)

 **Retrovir**
zidovudine (AZT, or ZDV)

 **Trizivir**
abacavir/lamivudine/zidovudine (ABC/3TC/AZT)

 **Videx EC**
didanosine (ddl)

 **Viracept**
nelfinavir (NFV)

 **Viramune XR**
nevirapine (NVP)

 **Zerit**
stavudine (d4T)

Walgreens Trusted si

SPONSOR OF THE 2019 POSITIVELY AWARE HIV DRUG CHART
For a list of Walgreens pharmacies that specialize in HIV/AIDS care and other resources, go to HIV.walgreens.com. For the complete 2019 POSITIVELY AWARE HIV Drug Guide, go to positivelyaware.com. To order more copies of the 2019 POSITIVELY AWARE HIV Drug Chart or HIV Drug Guide, email distribution@tpan.com.

© 2019 POSITIVELY AWARE